

University of Groningen

Circular polarization of radio emission from air showers in thunderstorm conditions

Trinh, T. N. G.; Scholten, O.; Bonardi, A.; Buitink, S.; Corstanje, A.; Ebert, U.; Enriquez, J. E.; Falcke, H.; Hörandel, J. R.; Mitra, P.

Published in:

7th International Conference on Acoustic and Radio EeV Neutrino Detection Activities

DOI:

[10.1051/epjconf/201713503002](https://doi.org/10.1051/epjconf/201713503002)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Trinh, T. N. G., Scholten, O., Bonardi, A., Buitink, S., Corstanje, A., Ebert, U., Enriquez, J. E., Falcke, H., Hörandel, J. R., Mitra, P., Mulrey, K., Nelles, A., Thoudam, S., Rachen, J. P., Rossetto, L., Rutjes, C., Schellart, P., ter Veen, S., & Winchen, T. (2017). Circular polarization of radio emission from air showers in thunderstorm conditions. In *7th International Conference on Acoustic and Radio EeV Neutrino Detection Activities: ARENA 2016* (Vol. 135). [3002] <https://doi.org/10.1051/epjconf/201713503002>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Circular polarization of radio emission from air showers in thunderstorm conditions

T.N.G.Trinh^{1,*}, O. Scholten^{1,2}, A. Bonardi³, S. Buitink⁴, A. Corstanje³, U. Ebert^{5,6}, J. E. Enriquez³, H. Falcke^{3,7,8,9}, J.R. Hörandel^{3,7}, P. Mitra⁴, K. Mulrey⁴, A. Nelles¹⁰, S. Thoudam¹¹, J.P. Rachen³, L. Rossetto³, C. Rutjes⁵, P. Schellart¹², S. ter Veen³, and T. Winchen⁴

¹KVI-Center for Advanced Radiation Technology, University Groningen, P.O. Box 72, 9700 AB Groningen, The Netherlands

²Interuniversity Institute for High-Energy, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussels, Belgium

³Department of Astrophysics/IMAPP, Radboud University Nijmegen, P.O. Box 9010, 6500 GL Nijmegen, The Netherlands

⁴Astrophysical Institute, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussels, Belgium

⁵Center for Mathematics and Computer Science (CWI), PO Box 94079, 1090 GB Amsterdam, The Netherlands

⁶Department of Applied Physics, Eindhoven University of Technology (TU/e), PO Box 513, 5600 MB Eindhoven, The Netherlands

⁷NIKHEF, Science Park Amsterdam, 1098 XG Amsterdam, The Netherlands

⁸Netherlands Institute of Radio Astronomy (ASTRON), Postbus 2, 7990 AA Dwingeloo, The Netherlands

⁹Max-Planck-Institut für Radioastronomie, P.O. Box 20 24, Bonn, Germany

¹⁰Department of Physics and Astronomy, University of California Irvine, Irvine, CA 92697-4575, USA

¹¹Department of Physics and Electrical Engineering, Linnéuniversitetet, 35195 Växjö, Sweden

¹²Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA

Abstract. We present measured radio emission from cosmic-ray-induced air showers under thunderstorm conditions. We observe for these events large differences in intensity, linear polarization and circular polarization from the events measured under fair-weather conditions. This can be explained by the effects of atmospheric electric fields in thunderclouds. Therefore, measuring the intensity and polarization of radio emission from cosmic ray extensive air showers during thunderstorm conditions provides a new tool to probe the atmospheric electric fields present in thunderclouds.

1 Introduction

Lightning initiation [1] and propagation [2] are driven by the electric fields in a thunderstorm. As it is very challenging to measure these fields in situ before or after the discharge and up to now essentially impossible during the discharge, there are ongoing controversies related to the physical mechanisms. A new non-intrusive method of field measurements is offered with LOFAR. It was discovered that under thunderstorm conditions, the intensity and polarizations of radio emission from

*e-mail: t.n.g.trinh@rug.nl

air showers differ considerably [3] from events in fair-weather conditions, so they can be used to probe the atmospheric electric fields present in thunderclouds.

Radio emission from air showers is measured by many experiments. Here we present the measurement with the LOw-Frequency ARay (LOFAR) radio telescope using the Low-Band Antennas (LBA). The LBA, having the frequency range of 10 - 90 MHz, are grouped into circular stations. The core of LOFAR, so-called 'Superterp', consists of 6 such stations, located in a ~ 320 m diameter region. The read-out of the LOFAR antennas for a cosmic-ray event is triggered by the LORA particle detector array at the Superterp. The measurements are done in both fair-weather and thunderstorm conditions.

2 Radio mechanism

The main mechanism for radio emission from air showers is the deflection of electrons and positrons in the showers in opposite direction by the Lorentz force exerted by the geomagnetic field [4, 5]. They form a transverse current and thus emit radiation polarized linearly along the direction of the Lorentz force, i.e. $\hat{\mathbf{e}}_{\mathbf{v} \times \mathbf{B}}$, where \mathbf{v} is the velocity of the shower and \mathbf{B} is the geomagnetic field.

A secondary contribution is from charge excess which is due to a build-up of negative charge in the shower front [6, 7]. This also produces a radio pulse polarized radially with respect to the shower symmetry axis. The signal observed on the ground is the superposition of the two contributions and thus depends on the viewing angle.

Under thunderstorm conditions, strong atmospheric electric fields affect the motion of shower particles and thus influence the radio emission from the showers. The electric field can be decomposed into two components

$$\mathbf{E} = \mathbf{E}_{\perp} + \mathbf{E}_{\parallel}, \quad (1)$$

where \mathbf{E}_{\perp} and \mathbf{E}_{\parallel} are perpendicular and parallel to the shower axis, respectively. Due to \mathbf{E}_{\perp} , there is a net force acting on particles

$$\mathbf{F} = q(\mathbf{E}_{\perp} + \mathbf{v} \times \mathbf{B}). \quad (2)$$

which gives rise to a change in both magnitude and direction of the transverse current. Thus, the magnitude and polarization of the radiation from the transverse current also change. Depending on the sign, the parallel component \mathbf{E}_{\parallel} may accelerate the particles and thus increase the number of particles at a certain energy. However, this effect is hardly seen in the signals observed by LOFAR LBA. These particles are trailing far behind the shower front, and therefore increase the intensity at lower frequencies [8]. Therefore, we assume that the parallel component vanishes in this work.

The complex voltages are expressed as $\varepsilon_j = E_j + i\hat{E}_j$ where E_j is sample j of the pulse and \hat{E}_j is its Hilbert transform. The Stokes parameters are given by

$$\begin{aligned} I &= \frac{1}{n} \sum_{j=0}^{n-1} (|\varepsilon_{j, \mathbf{v} \times \mathbf{B}}|^2 + |\varepsilon_{j, \mathbf{v} \times \mathbf{v} \times \mathbf{B}}|^2), \\ Q &= \frac{1}{n} \sum_{j=0}^{n-1} (|\varepsilon_{j, \mathbf{v} \times \mathbf{B}}|^2 - |\varepsilon_{j, \mathbf{v} \times \mathbf{v} \times \mathbf{B}}|^2), \\ U + iV &= \frac{2}{n} \sum_{j=0}^{n-1} (\varepsilon_{j, \mathbf{v} \times \mathbf{B}} \varepsilon_{j, \mathbf{v} \times \mathbf{v} \times \mathbf{B}}^*), \end{aligned} \quad (3)$$

In the LOFAR analysis, the summations are calculated for $n = 5$ samples of 5 ns each, centered around the peak of the pulse. Stokes I represents the intensity. The orientation of the linear polarization can be derived from Stokes Q and Stokes U. Stokes V is the amount of circular polarization.




Figure 1. Left panel: Linear polarization footprint of an air shower under thunderstorm condition. Each line represents the orientation of the linear polarization for each antenna. Right panel: Intensity footprint of an air shower under thunderstorm condition. The background is the result from CoREAS simulation. Each small circle displays the intensity for each antenna. There is a good agreement between the data and the simulation.

3 Fair-weather events vs thunderstorm events.

As discussed above, the events observed under thunderstorm conditions show large differences shown in the intensity and polarizations compared to fair-weather events.

3.1 Linear polarization

The angle of linear polarization is given by $\psi = \frac{1}{2} \tan^{-1}(\frac{U}{Q})$. In fair-weather events, the polarization over all antennas is mainly along the expected $\hat{e}_{v \times B}$ direction with some small deviation because of the contribution from charge excess. In thunderstorm events, the net force \mathbf{F} changes the direction of the transverse current and thus the orientation of the linear polarization is no longer along the $\hat{e}_{v \times B}$ direction, as it can be seen in the left panel of Fig. (1).

3.2 Intensity footprint

Unlike fair-weather events showing a bean-shaped intensity footprint due to the interference of the transverse current and charge excess components, the ones in thunderstorm conditions show a ring-like structure as seen in the right panel of Fig. (1). The observed structure for this event can be understood as the effect of an electric field that is build up in two layers. The upper layer starts at a height of 8 km above the ground and extends to 2.9 km with a strength of $|E_U| = 50$ kV/m. At the height of 2.9 km, the electric field changes such that the net force is inversed and its strength reduces to $|E_L| = 26.5$ kV/m. Two layers are needed to introduce the destructive interference between the emission from the two layers, which results in the ring-like structure in the intensity footprint.

3.3 Circular polarization

In fair-weather events, due to the time-shift between the pulses radiated from the transverse current component and the charge-excess component, there is a small amount of circular polarization. This




Figure 2. Circular polarization of an air shower under thunderstorm condition. The blue dots are simulated results and the red ones are data.

circular polarization vanishes at the core of the shower and depends on the azimuth angle of antennas [9]. However, in thunderstorm events we observe large amount of circular polarization near the core of the shower as shown in Fig. (2). This can only be explained by the change of the transverse current due to the change of the atmospheric electric field. Therefore, the full set of Stokes parameters helps to have a better understanding of the structure of atmospheric electric fields.

4 Conclusion

Using only the intensity footprint of the events measured under thunderstorm conditions, one can probe the atmospheric electric fields. It shows that we are sensitive to the height where the electric field changes and the relative strength between the fields in the two layers. The structure of electric fields can be reconstructed better by using both the intensity and the polarization information.

References

- [1] A. Dubinova et al., Phys. Rev. Lett. **115**, 015002 (2015)
- [2] J.R. Dwyer, M.A. Uman, Physics Reports **534**, 147 (2014), the Physics of Lightning
- [3] P. Schellart et al., Phys. Rev. Lett. **114**, 165001 (2015)
- [4] F.D. Kahn, I. Lerche, Royal Society of London Proceedings Series A **289**, 206 (1966)
- [5] O. Scholten, K. Werner, F. Rusydi, Astroparticle Physics **29**, 94 (2008)
- [6] G. Askaryan, J. Phys. Soc. Japan **Vol: 17, Suppl. A-III** (1962)
- [7] K.D. de Vries et al., Astroparticle Physics **34**, 267 (2010)
- [8] T.N.G. Trinh et al., Phys. Rev. D **93**, 023003 (2016)
- [9] O. Scholten et al. (2016), accepted by Phys. Rev. D (11/2016)